

Nuclear Receptor Coactivators: Regulators of Steroid Action in Brain and Behaviour

M. J. Tetel and K. D. Acharya

Neuroscience Program, Wellesley College, Wellesley, MA, USA.

Journal of Neuroendocrinology

Steroid hormones act in specific regions of the brain to alter behaviour and physiology. Although it has been well established that the bioavailability of the steroid and the expression of its receptor is critical for understanding steroid action in the brain, the importance of nuclear receptor coactivators in the brain is becoming more apparent. The present review focuses on the function of the p160 family of coactivators, which includes steroid receptor coactivator-1 (SRC-1), SRC-2 and SRC-3, in steroid receptor action in the brain. The expression, regulation and function of these coactivators in steroid-dependent gene expression in both brain and behaviour are discussed.

Correspondence to:

Marc J. Tetel, Neuroscience Program, Wellesley College, 106 Central Street, Wellesley, MA 02481, USA (e-mail: mtetel@wellesley.edu).

Key words: steroid receptor coactivator-1 (SRC-1/NcoA-1), SRC-2 (GRIP1/TIF2/NCoA-2), androgen receptor, oestrogen receptor, progesterin receptor, sex behaviour, hypothalamus, steroid hormones

doi: 10.1111/jne.12065

Steroid hormones act in the brain to profoundly influence behaviour and physiology. These hormones elicit many of their biological effects by binding to their respective receptors, which are members of the steroid/nuclear receptor superfamily of transcriptional activators. Receptors for oestrogens (ER), progestins (PR) and androgens (AR) can act in a classic genomic mechanism of action by binding directly to target DNA to alter transcription (1). In addition, these receptors can act in a rapid, nonclassical manner, involving receptors located at the membrane that activate intracellular signalling pathways in the brain (2–7). In the classic genomic mechanism of action, nuclear receptor coregulators enhance (coactivators) or repress (corepressors) the transcriptional activity of steroid receptors. Over 350 coregulators have been identified to function with the large superfamily of nuclear receptors (8). Knowledge of the function of these coregulators in behaviour, physiology and disease is growing rapidly. The present review focuses on the role of the p160 steroid receptor coactivator family in the classic genomic mechanism of ER and PR action in the brain and the regulation of behaviour.

Genomic mechanisms of ER and PR action

ER and PR, as well as other steroid receptors, have a modular domain structure consisting of a highly variable amino-terminal region (N-domain), a conserved central DNA-binding domain and a

carboxy-terminal ligand binding domain (1,9). In general, steroid receptors have two transcriptional activation domains in the amino (AF-1) and carboxyl (AF-2) termini (10). Intracellular ER are expressed as two subtypes, α and β , which are transcribed from different genes (11,12). These subtypes differ in their abilities to bind different ligands and regulate transcription (13,14), distribution in the brain (15–18) and regulation of behaviour (19–23). In addition, there are ER splice variants (24–26) that bind differentially with coactivators (27) and may provide another level of regulation. In most species, PR are expressed in two forms, the full-length PR-B and the N-terminally truncated PR-A, which are encoded by the same gene but are under the regulation of alternate promoters and internal translation start sites (28,29). Under certain cell and promoter contexts, human PR-B is a stronger transcriptional activator than PR-A (30–32) and PR-A can repress the transcriptional activity of PR-B. These differences are most likely the result of an additional AF domain in the N-terminus of PR-B (33) and a transcriptional inhibitory region that has been identified in PR-A (32,34), respectively. In further support of these differences in PR-A and PR-B, these two PR isoforms appear to have distinct functions in reproductive behaviour and physiology (35,36).

In the classic, ligand-dependent, genomic mechanism of action of ER, PR and other steroid receptors, they are complexed with several chaperone molecules, including heat shock proteins (hsp), in the absence of hormone. Upon binding hormone, receptors undergo

a conformational change that causes the dissociation of hsp, allowing receptors to dimerise (37). Activated receptors bind directly to specific steroid response elements (SRE) and SRE-like sequences in the promoter regions of target genes (1,9). Binding of receptors to DNA increases or decreases gene transcription by altering the rate of recruitment of general transcription factors and influencing the recruitment of RNA polymerase II to the initiation site (38,39). It is generally considered that oestrogens and progestins can act in the brain via their respective receptors to alter neuronal gene transcription in a fashion similar to that described above, resulting in profound changes in behaviour and physiology (4,40,41).

Nuclear receptor coregulators

Coregulators consist of coactivators and corepressors that are required for efficient transcriptional regulation by nuclear receptors (8,42). Corepressors and their complexes associate with nuclear receptors when unliganded or bound to antagonists and serve to repress nuclear receptor transcription by recruiting corepressor complexes to the cis-active elements in the promoter and enhancers of target genes (42). Nuclear receptor coactivators, which are the focus of the present review, dramatically enhance the transcriptional activity of ER and PR, as well as other nuclear receptors, by acting as bridging molecules between the receptor and the general transcription machinery and modifying chromatin within the promoter and enhancer regions by histone acetylation, methylation and phosphorylation (42,43). Under most conditions, steroid receptors interact with coactivators in the presence of an agonist but not in the absence of ligand or in the presence of an antagonist or a selective receptor modulator (44–47); but see also additional studies (48–50). *In vitro* studies indicate that recruitment of nuclear receptor coactivators is rate-limiting in steroid receptor-mediated gene transcription (42,51). In further support of the importance of nuclear receptor coactivators in steroid-dependent transcription *in vitro*, squelching, or the repression of the transcriptional activity of one steroid receptor by another, is reversed by the addition of coactivators (44). Thus, a critical component of efficient steroid dependent transcription is the recruitment by receptors of nuclear receptor coactivators to the complex (8,42). Finally, the significance of both coactivators and corepressors in a variety of diseases, including hormone-dependent cancer and some neurological disorders, is becoming more apparent (8).

The p160 steroid receptor coactivator (SRC) family

The SRC family of p160 proteins consists of SRC-1 (NcoA-1), SRC-2 (GRIP1/TIF2/NCoA-2) and SRC-3 (AIB1/TRAM-1/ACTR/RAC3/pCIP). Nuclear receptor coactivators, including the SRC coactivator family, share a general set of characteristics. The SRC family of coactivators physically interacts with steroid receptors, including ER, PR, AR and receptors for glucocorticoids, in a ligand-dependent manner (43). The SRCs physically interact with agonist-bound receptors through centrally-located multiple LXXLL motifs (L, leucine; X, any amino acid) that make up nuclear receptor boxes. The SRCs and other coactivators do not bind DNA and thus distinguish them from

traditional transcription factors. The C-terminus of the SRCs contains two activation domains: AD-1 and AD-2. The N-terminus contains a third activation domain (AD-3) and a basic helix loop helix-Per Arnt Sims (bHLH-PAS) motif, which is the most conserved domain within this family of proteins. The activation domains interact with secondary coactivators known as co-coactivators (43). These co-coactivators modify chromatin to facilitate binding of regulatory proteins and general transcription factors.

The p160 SRC family in reproductive physiology and behaviour

Expression and regulation in the brain

Sex steroids, including oestrogens, progestins and androgens, are required for brain development and reproductive behaviour in rodents and birds. Therefore, both rodents and birds represent excellent models for studying coactivator function in the brain. In male and female rodents, SRC-1 mRNA and protein are expressed at high levels in the cortex, hippocampus, cerebellum and hypothalamus (52–58,59: 94). In addition, the SRC-1 isoform, full length SRC-1a, is found in high levels in the rodent hypothalamus, whereas levels of the C-terminally truncated SRC-1e are higher in the nucleus accumbens, thalamus and amygdala (54). Recently, this SRC-1a : SRC-1e ratio has been shifted in the central nucleus of the amygdala using antisense targeting the SRC-1e isoform, which may prove valuable for studying the functions of these SRC isoforms in the brain (60). The expression of SRC-1 in the female rat brain appears to decline as the animal ages, suggesting a loss of steroid sensitivity (61). SRC-2 is highly expressed throughout the hippocampus, amygdala and hypothalamus, including the medial preoptic area (MPOA), ventral medial nucleus (VMN), arcuate nucleus (ARC), bed nucleus of the stria terminalis, supraoptic nucleus and suprachiasmatic nucleus (58,62–64). Although it is not known whether a sex difference exists, SRC-3 is expressed predominantly in the hippocampus and very sparsely in the hypothalamus in both male and female rodents (58,64).

The avian brain provides an excellent model for studying steroid action since singing and nonsinging birds respond to steroids. Songbirds have a specific group of interconnected nuclei called the song control system that are required for singing and are sexually dimorphic and steroid-sensitive (65). In the songbird zebra finch, AR and ER are expressed in the song control nuclei from early post-hatching ages (66–68). Injection of 17β -oestradiol in early post-hatching females masculinises the song system and makes the females capable of singing as adults (69–72). Although the organisational effects of oestrogens are limited to early development in zebra finches, in other songbirds, such as canaries, manipulation of these hormones during adulthood affects the size of song nuclei and song (73). In the quail, a nonsinging bird, steroids regulate both appetitive and consummatory (copulatory) male sexual behaviour (74–77). In males, AR and ER are expressed in the medial preoptic nucleus (POM) of quail and are required for both aspects of sexual behaviour (74,78). Members of the p160 family of steroid receptor coactivators, SRC-1 and SRC-2, are expressed in both

songbirds and nonsinging birds. SRC-1 mRNA is expressed as early as post-hatching day 1 in the telencephalon of zebra finches, and in the song control nuclei and hypothalami of adult canaries and zebra finches (79,80). Interestingly, SRC-1 mRNA and protein show a male-biased expression in the song nucleus HVC of adult canaries and zebra finches, respectively (79,80). In the quail brain, SRC-1 is expressed in the steroid sensitive areas, including the POM and bed nucleus of stria terminalis (79). Similar to SRC-1, quail POM expresses SRC-2 protein in a level that is similar in males and females (81).

For coactivators to function with steroid receptors, they must be expressed in the same cells. Oestradiol-priming dramatically increases the expression of PR in a variety of rodent brain regions, including the MPOA, VMN, ARC and the midbrain central grey (82–87). We found that SRC-1 and SRC-2 are expressed in the majority of oestradiol-induced PR cells in regions involved in female reproduction, including the VMN, MPOA and ARC in rats and mice (62,88,89). Given that almost all oestradiol-induced PR cells in the hypothalamus contain ER α (84,85), these findings suggest that these coexpressing cells represent functional sites of interaction between steroid receptors and coactivators in the brain (62,88,89). In further support, SRC-1 was found to be expressed in oestrogen-sensitive pro-opiomelanocortin and steroidogenic factor-1 neurones in the arcuate nucleus and ventromedial hypothalamus (VMH), respectively (90).

It is assumed that coactivators are modulators of cellular responsiveness to steroids. In support, SRC-1 knockout mice, although fertile, have decreased responsiveness in progestin target tissues (91) and partial resistance to thyroid hormone (92). It is important to note that, in these mice, SRC-2 is up-regulated in steroid sensitive tissues, including the brain and testes, suggesting that increased expression of SRC-2 compensates for the loss of SRC-1 (91). Therefore, studying the regulation of coactivator expression is essential for understanding hormone action in the brain. A number of studies indicate that hormones can regulate coactivator expression in rodent and bird brains. In rodents, SRC-1 is expressed in a sexually dimorphic manner in the pituitary gland, with males having higher mRNA (52) and protein (93) levels than females. In further support, male rodents have higher levels of SRC-1 than females in a number of brain regions, including the dorsomedial hypothalamus, VMH and paraventricular nucleus (94). Ovariectomy decreases SRC-1 expression in the VMH, whereas oestradiol reverses this effect (95). In the hypothalamus of cycling female rats, SRC-1 levels were lowest during di-oestrus, and highest at pro-oestrus and oestrus, suggesting that ovarian hormones up-regulate SRC-1 (96). By contrast, ovariectomy did not alter SRC-1 levels in the hippocampus, suggesting that ovarian hormones do not regulate SRC-1 expression in this brain region (97). Interestingly, the endocrine disruptor 4-methylbenzylidene camphor (4-MBC), which has oestrogenic activity and impairs the thyroid axis, increases SRC-1 mRNA in the VMH and MPOA of female rats (98). Exposure to another endocrine disruptor, 3-benzylidene camphor (3-BC), during early development through adulthood increases SRC-1 mRNA levels in the MPOA of both males and females (99). These effects of 4-MBC and 3-BC on SRC-1 could enhance their oestrogenic effects and alter other nuclear receptor signalling pathways. Testosterone treatment does not alter SRC-1 expression in the

MPOA, bed nucleus of the stria terminalis (BNST), ARC and amygdala of castrated hamsters (100). However, testosterone decreases SRC-2 expression in the hypothalamus of male rats (63). Finally, thyroid hormone decreases SRC-1 expression in rat cortex and dentate gyrus (101) and neonatal mouse cerebellum (102). In adult birds, testosterone increases SRC-1 expression in the quail hypothalamus (103), whereas the administration of oestradiol, testosterone or aromatase inhibitor has no effect on SRC-1 expression in zebra finches (80).

In addition to gonadal steroids, it appears that glucocorticoids and stress can influence SRC-1 expression. Treatment of male rats with the synthetic glucocorticoid, dexamethasone, reduces SRC-1 mRNA in the brain but does not affect the other members of the p160 family of coactivators, SRC-2 and SRC-3 (104). In further support, adrenalectomised male rats exposed to high levels of corticosterone have decreased SRC-1e mRNA levels in the anterior pituitary but, interestingly, no changes were detected in the hippocampus (105). In rats, acute restraint stress decreases SRC-1 expression in the male and female hypothalamus and male frontal cortex, and increases SRC-1 levels in the male pituitary and the female hippocampus (93). Taken together, these studies suggest that glucocorticoids and stress may alter brain function by influencing coactivator expression in a brain region- and sex-specific manner.

Daylength has profound effects on reproduction and other neuroendocrine events (106). In male Siberian hamsters exposed to short days, we found reduced SRC-1 expression in the posteromedial BNST and posterodorsal medial amygdala (100). In addition, SRC-1 expression in the hippocampus, hindbrain and optic lobes change through the day in Japanese quail (103,107). Given that both Siberian hamsters and Japanese quail have seasonal cycles, these findings suggest that this photoperiodic regulation of SRC-1 contributes to androgen regulation of seasonal reproduction.

An increasing number of novel functions are being attributed to the p160 family of coactivators. For example, SRC-1 is predominantly expressed in neuronal lineage cell lines during neural stem cell differentiation (108). In addition, this expression of SRC-1 is higher in mature neurones than immature neurones, suggesting a role for SRC-1 in the differentiation of neural stem cells (108). Further investigation of coactivator expression will be essential to fully understand their function in hormone action.

In addition to regulation of coactivator expression, functional interaction of coactivators with receptors can be affected by post-translational modifications, such as phosphorylation, methylation and acetylation of coactivators (109). For example, SRC-1, SRC-2 and SRC-3 undergo phosphorylation at different sites (110–114), which can alter the conformation, stability and activity of these proteins (109,110). Given that these posttranslational modifications have been studied in cell culture systems, it will be important in future studies to explore whether these modifications occur in the brain and impact behaviour.

Regulation of steroid-dependent gene expression in the brain by coactivators

A classic example of steroid-dependent gene expression is the oestradiol-induction of PR in a variety of oestrogen-responsive tissues,

including the brain, breast and uterus (82–87). Induction of PR expression by oestradiol in the VMH is important for steroid-dependent female sexual behaviour in rodents (115). Therefore, we tested the hypothesis that SRC-1, along with the co-coactivator CREB-binding protein (CBP), is critical for modulating ER-mediated transactivation of the PR gene in the VMN. Infusions of antisense to SRC-1 and CBP mRNA into the VMN of adult female rats reduced the expression of ER-mediated activation of PR gene expression compared to controls (56). These findings extend previous *in vitro* studies indicating that SRC-1 and CBP act together to modulate ER and PR function (116,117). Another study in rodent brain supports these findings with respect to SRC-1 function in the ER-mediated induction of PR in the VMN, extending them to include a role for SRC-2 but not SRC-3 (64). In a mouse hypothalamic neuronal cell line, ER β bound to the ER β agonist, 3 β -diol, increased oxytocin gene mRNA levels and the occupancy of the oxytocin gene promoter by SRC-1 and CBP (118). These results suggest that SRC-1 and CBP form a coactivator complex that regulates oxytocin gene expression (118) and support the findings reported above showing that SRC-1 and CBP function in ER-mediated induction of PR in the brain (56).

In male quails, the volume of the POM, a critical brain region in male sexual behaviour, and aromatase expression is increased by testosterone treatment within 14 and 2 days, respectively (74). Interestingly, knocking down SRC-1 by antisense decreases the testosterone-dependent POM volume and aromatase immunoreactivity in male quails, suggesting a role for SRC-1 in testosterone-induced changes in brain structure and gene expression in birds (119). Although not a member of the p160 family of coactivators, another steroid receptor coactivator, ribosomal protein L7 (RPL7, aka L7/SPA), has been well-studied in the bird brain. RPL7 is part of the ribosomal complex required for transcription and translation (120) and has been shown to be a coactivator for ER α , PR and vitamin D receptor (121,122). In the song system of zebra finches, RPL7 protein shows a greater expression at posthatch day 1 and in adult males compared to females (123). Antisense administration to RPL7 mRNA increased neuronal death in HVC and Area X, suggesting a role for this coactivator in neuroprotection (124). Similar effects of reducing RPL7 were observed in neuronal cultures from posthatch day 1 males and females, with neuronal loss being greater in males compared to females. Oestradiol treatment prevented the neuronal loss caused by antisense to RPL7, suggesting that the neuroprotective effects of oestradiol are not dependent on ER α in this model (124,125).

In further support of a role for the p160 family of coactivators in modulating ER action in the brain, studies have recently been performed in human astrocytoma cell lines. Oestradiol treatment increases the number of cells in two (U373 and D54) astrocytoma cell lines (126). This effect appears to be mediated by ER α , given that the ER α agonist (propyl-pyrazole-triol; PPT), but not the ER β agonist (diarylpropionitrile), mimicked the effects of oestradiol on cell proliferation. Interestingly, coactivator silencing by RNA interference of SRC-1 (but not SRC-3) blocked the PPT-induced increase in cell number, suggesting that SRC-1 regulates the ER α -mediated increase in cell number in these astrocytoma cell lines (126). In a

related study, progesterone increases vascular endothelial growth factor expression (VEGF) in this D54 astrocytoma cell line (127). Silencing of SRC-1 reduced VEGF protein levels following progesterone treatment, suggesting that SRC-1 is important in modulating the expression of this progesterone sensitive gene (127). Future studies in the brain and cell lines will be critical for further determining the function of coactivators in modulating steroid action in the brain.

Coactivators modulate steroid-dependent behaviours

Given that nuclear receptor coactivators appear to be essential for hormone-dependent gene expression in the brain, we tested the hypothesis that coactivators act in the brain to modulate the expression of hormone-dependent behaviours (56,128). Female rats treated with antisense to both SRC-1 and CBP mRNA into the VMN showed lower levels of steroid-dependent lordosis compared to scrambled-treated controls (56). Another study supported these findings with SRC-1 and extended them to include a role for SRC-2 in hormone-dependent lordosis (64). In further support of the gene expression studies discussed above, SRC-3 did not appear to function in the brain in steroid-dependent lordosis (64). Given that ER α (and not ER β) appears to mediate female sexual behaviour in rats (129), these findings suggest that SRC-1 and SRC-2 are functioning with ER α to elicit these effects on behaviour.

One limitation of the behavioural experiments discussed above is that they do not isolate the effects of coactivators on specific ER- and PR-dependent aspects of female sexual behaviour. Therefore, we designed experiments aiming to investigate whether coactivators act specifically with ER or PR in the brain to influence behaviour in rats (128). To test the hypothesis that coactivators modulate ER-mediated aspects of female sexual behaviour, animals were injected with two slightly higher doses of oestradiol alone, which elicit lordosis (41). Antisense to SRC-1 and CBP infused into the VMN of animals treated with oestradiol decreased the frequency and intensity of lordosis, suggesting that these coactivators modulate ER-mediated aspects of female sexual behaviour (128). To investigate whether coactivators act with PR in the brain to influence behaviour, we took advantage of the fact that proceptive behaviours by the female, such as ear-wiggling and hopping and darting that serve to solicit interaction by the male, are PR-dependent (130,131). In this experiment, antisense to SRC-1 and CBP mRNA was infused into the VMN after priming with oestradiol around the time of progesterone administration. This timing of coactivator antisense infusion allowed for the disruption of PR activity but did not alter induction of PR by oestradiol. Females treated with antisense to coactivators had a reduced frequency of PR-dependent ear-wiggling and hopping and darting but not PR-dependent receptivity (128). These findings suggest that a reduction of SRC-1 by antisense disrupted the activity of PR signalling pathway(s) influencing proceptivity, whereas alternate PR signalling pathways that regulate PR-dependent receptivity remained intact and functional. Thus, it appears that coactivators function in the brain to modulate both PR- and ER-specific aspects of steroid-dependent female sexual behaviours in rodents.

Studies in male quails provide further support for a role of SRC-1 and SRC-2 in regulating behaviour. Antisense to SRC-1 in the POM of quail inhibited both AR- and ER-mediated sexual behaviour (132). In quails, strutting and crowing by males as responses towards females are androgen dependent, whereas mount attempts, mounts and cloacal contact movements by the male are oestrogen-dependent (76,77). Testosterone injection induces these behaviours by directly acting on AR and on ER following the aromatisation of testosterone to oestrogens. Antisense to SRC-1 blocked all of these testosterone-mediated male sexual behaviours, which were reinstated after terminating the antisense treatment (132). SRC-2 is also required in reproductive behaviour, as demonstrated by a reduction in the size of the POM, as well as a decrease in testosterone-induced male sexual behaviour, following SRC-2 antisense injection into the third ventricle (81).

Coactivators from brain associate with ER and PR

As noted above, one of the criteria of nuclear receptor coactivators is that they physically associate with receptors. To test the hypotheses that members of the p160 family of steroid receptor coactivators from brain physically associate with ER and PR subtypes in a ligand-dependent manner, we developed pull-down assays with brain tissue from female rodents.

SRC-1 from rat hypothalamic or hippocampal extracts interacted with Flag-tagged ER α and ER β when bound to oestradiol, which was confirmed by mass spectrometry (133). Little to no association of SRC-1 from brain with ER α or ER β was detected in the absence of ligand or in the presence of tamoxifen, a selective ER modulator (SERM). These findings suggest that SRC-1 from brain interact with ER in a ligand-dependent manner and that the SERM tamoxifen functions as an antagonist in this assay to prevent receptor-coactivator interactions. In further support, the ER α agonist, PPT, promoted physical association between ER α and SRC-1 in the hypothalamus, as detected by co-immunoprecipitation (90). These results support our previous findings that SRC-1 action in the hypothalamus is important for maximal ER-mediated transactivation of the PR gene and expression of female sexual behaviour (56,128). SRC-1 may function with both ER subtypes in the hippocampus to differentially modulate the effects of oestrogens on cognition and stress (19,23,134–137). Interestingly, SRC-1 from the hippocampus interacted equally with ER α and ER β , whereas SRC-1 obtained from hypothalamic extracts interacted more with ER α than with ER β , suggesting that other cofactors involved in these protein-protein interactions have different expression patterns in these brain regions. In addition, it is possible that SRC-1 undergoes distinct post-translational modifications (e.g. phosphorylation) in these two brain regions, leading to differential interactions with receptors.

Similar to findings with SRC-1 and as also confirmed by mass spectrometry, SRC-2 from hypothalamus or hippocampus interacted with ER α in a ligand-dependent manner (62). However, in dramatic contrast to SRC-1, SRC-2 from the brain showed little to no interactions with ER β under any ligand conditions. This weak association of oestradiol-bound ER β with SRC-2 from brain is in contrast to

cell culture studies indicating that over-expressed SRC-2 interacts with ER β (138–141). It is possible that the over-expression of coactivators leads to altered interactions with receptors and/or that the presence of other factors in the brain may mediate appropriate receptor-coactivator associations. Taken together, these findings suggest that it is important to use biologically-relevant tissue when studying these receptor-coactivator interactions. Finally, these differential interactions between SRC-2 and ER α and ER β may contribute to the functional differences of these ER subtypes in the brain (19). In future studies, it will be important to explore the possibilities that coactivators, including the SRCs, function in non-genomic oestrogen signalling pathways in the brain.

Interactions between coactivators from brain and the PR isoforms have also been studied. SRC-1 from rat hypothalamic or hippocampal extracts interacted with both GST-tagged PR-A and PR-B when bound to the agonist R5020 but not in the absence of ligand or in the presence of the selective PR modulator, RU486 (133). These agonist-dependent interactions between PR and SRC-1 from brain support our previous work indicating a role for hypothalamic SRC-1 in PR-dependent female sexual behaviour (128) and provide evidence that SRC-1 may contribute to progestin effects in the hippocampus on memory (142,143). Interestingly, we found that SRC-1 from the hypothalamus or hippocampus interacts more with PR-B, than with PR-A. In regard to SRC-2, we found that this coactivator interacted with PR-B (but not PR-A) in a ligand-dependent manner. Furthermore, cell culture studies suggest that, under certain circumstances, human PR-B is a stronger transcriptional activator than PR-A (32,144,145), likely as a result of the additional activation function (AF-3) of PR-B (33,146). Our findings showing that these coactivators interact more with PR-B than PR-A are consistent with some cell culture studies (145) and suggest a mechanism by which PR-B may be a stronger transcriptional activator than PR-A. However, it should be noted that, although studies using PR-A and PR-B specific knockouts reveal that both receptors are important for the full display of progesterone-facilitated lordosis, PR-A has a greater role than PR-B in ligand-independent lordosis facilitated by the cyclic AMP analogue, 8-bromo-cAMP (35). We are currently using mouse PR and mouse brain tissue to explore PR-coactivator interactions. In future studies, it will be important to investigate the function of the SRCs and other coactivators in the ligand-independent activation of PR in the rodent brain. Understanding how nuclear receptor coactivators interact with various steroid receptors, and their subtypes, is critical for understanding how hormones act in different brain regions to profoundly influence physiology and behaviour. Ultimately, mass spectrometry analyses of these receptor-coactivator interactions using brain tissue may allow the identification of novel coregulators involved in the steroid receptor complex in the brain.

Conclusions

Following the discovery of the p160 family of coactivators and other nuclear receptor coactivators, findings from *in vitro* and cell culture studies have revealed much about the function of coactivators in steroid action. More recently, approaches using animal




Fig. 1. Diagram depicting the concept that steroid receptors (SR) recruit different sets of nuclear receptor coactivators (NRC) to enhance transactivation of distinct target genes and elicit different behaviours.

models and neuronal cell lines have greatly expanded our knowledge of coactivator function and enabled us to better understand how these coactivators modulate steroid action in brain and influence complex behaviours. In addition, our recent receptor–coactivator interaction studies using rodent brains discussed above highlight the significance of using biologically-relevant tissue for exploring these important interactions. A critical question in neuroendocrinology is how individual cells respond to steroids and how this responsiveness can change over time or with experience. The regulation and expression of a large diversity of nuclear receptor coactivators, including the p160 family of coactivators, provides a mechanism by which individual cells in specific brain regions can differentially respond to hormones and enable the adjustment of this sensitivity to steroids in response to changes in external stimuli. In addition, recruitment of different members of the p160 family of coactivators by receptors may lead to distinct signalling pathways and behaviours (Fig. 1). In support, *in vitro* studies show that ERs recruit either SRC-1 or SRC-2 depending on the oestrogen response element (147). Future research using a variety of animal models, including rodent and bird models, will continue to investigate the function of these important regulatory proteins in behaviour, physiology and disease.

Acknowledgements

Studies contributed by the authors' laboratory were supported by grants received from the National Institutes of Health R01 DK61935 (M.J.T.).

Received 15 April 2013,
revised 3 June 2013,
accepted 18 June 2013

References

- Mangelsdorf DJ, Thummel C, Beato M, Herrlich P, Schütz G, Umesono K, Blumberg B, Kastner P, Mark M, Chambon P, Evans RM. The nuclear receptor superfamily: the second decade. *Cell* 1995; **83**: 835–839.
- Cornil CA, Ball GF, Balthazart J. Rapid control of male typical behaviours by brain-derived estrogens. *Front Neuroendocrinol* 2012; **33**: 425–446.
- Vasudevan N, Pfaff DW. Membrane-initiated actions of estrogens in neuroendocrinology: emerging principles. *Endocr Rev* 2007; **28**: 1–19.
- Tetel MJ, Lange CA. Molecular genomics of progestin actions. In: Pfaff DW, Arnold AP, Etgen AM, Fahrbach SE, Rubin RT, eds. *Hormones, Brain and Behavior*. San Diego, CA: Academic Press, 2009: 1439–1465.
- Mani SK, Mermelstein PG, Tetel MJ, Anesetti G. Convergence of multiple mechanisms of steroid hormone action. *Horm Metab Res* 2012; **44**: 569–576.
- Kelly MJ, Ronnekleiv OK. Membrane-initiated actions of estradiol that regulate reproduction, energy balance and body temperature. *Front Neuroendocrinol* 2012; **33**: 376–387.
- Micevych PE, Dewing P. Membrane-initiated estradiol signaling regulating sexual receptivity. *Front Endocrinol* 2011; **2**: 26.
- Lonard DM, O'Malley BW. Nuclear receptor coregulators: modulators of pathology and therapeutic targets. *Nat Rev Endocrinol* 2013; **8**: 598–604.
- Tsai MJ, O'Malley BW. Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu Rev Biochem* 1994; **63**: 451–486.
- Tora L, White J, Brou C, Tasset D, Webster N, Scheer E, Chambon P. The human estrogen receptor has two independent non-acidic transcriptional activation functions. *Cell* 1989; **59**: 477–487.
- Jensen EV, Suzuki T, Kawasima T, Stumpf WE, Jungblut PW, de Sombre ER. A two-step mechanism for the interaction of estradiol with rat uterus. *Proc Natl Acad Sci USA* 1968; **59**: 632–638.
- Kuiper GGJM, Enmark E, Peltö-Huikko M, Nilsson S, Gustafsson J. Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci USA* 1996; **93**: 5925–5930.
- Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Häggblad J, Nilsson S, Gustafsson J. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* 1997; **138**: 863–870.
- Delaunay F, Pettersson K, Tujague M, Gustafsson JA. Functional differences between the amino-terminal domains of estrogen receptors alpha and beta. *Mol Pharmacol* 2000; **58**: 584–590.
- Shughrue PJ, Lane MV, Merchenthaler I. Comparative distribution of estrogen receptor-alpha and -beta mRNA in the rat central nervous system. *J Comp Neurol* 1997; **388**: 507–525.
- Greco B, Allegretto EA, Tetel MJ, Blaustein JD. Coexpression of ER beta with ER alpha and progestin receptor proteins in the female rat forebrain: effects of estradiol treatment. *Endocrinology* 2001; **142**: 5172–5181.
- Osterlund M, Kuiper GG, Gustafsson JA, Hurd YL. Differential distribution and regulation of estrogen receptor-alpha and -beta mRNA within the female rat brain. *Brain Res Mol Brain Res* 1998; **54**: 175–180.
- Mitra SW, Hoskin E, Yudkovitz J, Pear L, Wilkinson HA, Hayashi S, Pfaff DW, Ogawa S, Rohrer SP, Schaeffer JM, McEwen BS, Alves SE. Immunolocalization of estrogen receptor beta in the mouse brain: comparison with estrogen receptor alpha. *Endocrinology* 2003; **144**: 2055–2067.
- Tetel MJ, Pfaff DW. Contributions of estrogen receptor-alpha and estrogen receptor-beta to the regulation of behavior. *Biochim Biophys Acta* 2010; **1800**: 1084–1089.
- Ogawa S, Eng V, Taylor J, Lubahn DB, Korach KS, Pfaff DW. Roles of estrogen receptor-alpha gene expression in reproduction-related behaviors in female mice. *Endocrinology* 1998; **139**: 5070–5081.
- Musatov S, Chen W, Pfaff DW, Kaplitt MG, Ogawa S. RNAi-mediated silencing of estrogen receptor {alpha} in the ventromedial nucleus of hypothalamus abolishes female sexual behaviors. *Proc Natl Acad Sci USA* 2006; **103**: 10456–10460.
- Ogawa S, Chan J, Chester AE, Gustafsson JA, Korach KS, Pfaff DW. Survival of reproductive behaviors in estrogen receptor beta gene-deficient (betaERKO) male and female mice. *Proc Natl Acad Sci USA* 1999; **96**: 12887–12892.
- Bodo C, Rissman EF. New roles for estrogen receptor beta in behavior and neuroendocrinology. *Front Neuroendocrinol* 2006; **27**: 217–232.

- 24 Koehorst SGA, Cox JJ, Donker GH, Lopes da Silva S, Burbach JPH, Thijsen JHH, Blankenstein MA. Functional analysis of an alternatively spliced estrogen receptor lacking exon 4 isolated from MCF-7 breast cancer cells and meningioma tissue. *Mol Cell Endocrinol* 1994; **101**: 237–245.
- 25 Madsen MW, Reiter BE, Lykkesfeldt AE. Differential expression of estrogen receptor mRNA splice variants in the tamoxifen resistant human breast cancer cell line, MCF-7/TAMR-1 compared to the parental MCF-7 cell line. *Mol Cell Endocrinol* 1995; **109**: 197–207.
- 26 Ishunina TA, Swaab DF. Hippocampal estrogen receptor- α splice variant TADD1 in the human brain in aging and Alzheimer's disease. *Neuroendocrinology* 2009; **89**: 187–199.
- 27 Bollig A, Miksicek RJ. An estrogen receptor- α splicing variant mediates both positive and negative effects on gene transcription. *Mol Endocrinol* 2000; **14**: 634–649.
- 28 Kastner P, Krust A, Turcotte B, Stropp U, Tora L, Gronemeyer H, Chambon P. Two distinct estrogen-regulated promoters generate transcripts encoding the two functionally different human progesterone receptor forms A and B. *EMBO J* 1990; **9**: 603–614.
- 29 Horwitz KB, Sheridan PL, Wei LL, Krett NL. Human progesterone receptors: synthesis, structure, and phosphorylation. *Prog Clin Biol Res* 1990; **32**: 241–252.
- 30 Vegeto E, Shahbaz MM, Wen DX, Goldman ME, O'Malley BW, McDonnell DP. Human progesterone receptor A form is a cell- and promoter-specific repressor of human progesterone receptor B function. *Mol Endocrinol* 1993; **7**: 1244–1255.
- 31 Tung L, Kamel Mohamed M, Hoeffler JP, Takimoto GS, Horwitz KB. Antagonist-occupied human progesterone B-receptors activate transcription without binding to progesterone response elements and are dominantly inhibited by A-receptors. *Mol Endocrinol* 1993; **7**: 1256–1265.
- 32 Giangrande PH, Pollio G, McDonnell DP. Mapping and characterization of the functional domains responsible for the differential activity of the A and B isoforms of the human progesterone receptor. *J Biol Chem* 1997; **272**: 32889–32900.
- 33 Sartorius CA, Melville MY, Hovland AR, Tung L, Takimoto GS, Horwitz KB. A third transactivation function (AF3) of human progesterone receptors located in the unique N-terminal segment of the B-isoform. *Mol Endocrinol* 1994; **8**: 1347–1360.
- 34 Hovland AR, Powell RL, Takimoto GS, Tung L, Horwitz KB. An N-terminal inhibitory function, IF, suppresses transcription by the A-isoform but not the B-isoform of human progesterone receptors. *J Biol Chem* 1998; **273**: 5455–5460.
- 35 Mani SK, Reyna AM, Chen JZ, Mulac-Jericevic B, Conneely OM. Differential response of progesterone receptor isoforms in hormone-dependent and -independent facilitation of female sexual receptivity. *Mol Endocrinol* 2006; **20**: 1322–1332.
- 36 Mulac-Jericevic B, Conneely OM. Reproductive tissue selective actions of progesterone receptors. *Reproduction* 2004; **128**: 139–146.
- 37 DeMarzo A, Beck CA, Oñate SA, Edwards DP. Dimerization of mammalian progesterone receptors occurs in the absence of DNA and is related to the release of the 90-kDa heat shock protein. *Proc Natl Acad Sci USA* 1991; **88**: 72–76.
- 38 Klein-Hitpass L, Tsai SY, Weigel NL, Allan GF, Riley D, Rodriguez R, Schrader WT, Tsai MJ, O'Malley BW. The progesterone receptor stimulates cell-free transcription by enhancing the formation of a stable preinitiation complex. *Cell* 1990; **60**: 247–257.
- 39 Kininis M, Chen BS, Diehl AG, Isaacs GD, Zhang T, Siepel AC, Clark AG, Kraus WL. Genomic analyses of transcription factor binding, histone acetylation, and gene expression reveal mechanistically distinct classes of estrogen-regulated promoters. *Mol Cell Biol* 2007; **27**: 5090–5104.
- 40 Pfaff D. Hormone-driven mechanisms in the central nervous system facilitate the analysis of mammalian behaviours. *J Endocrinol* 2005; **184**: 447–453.
- 41 Blaustein JD, Mani SK. Feminine sexual behavior from neuroendocrine and molecular neurobiological perspectives. In: Blaustein JD, ed. *Handbook of Neurochemistry and Molecular Neurobiology*. New York, NY: Springer, 2006: 95–150.
- 42 Rosenfeld MG, Lunyak VV, Glass CK. Sensors and signals: a coactivator/corepressor/epigenetic code for integrating signal-dependent programs of transcriptional response. *Genes Dev* 2006; **20**: 1405–1428.
- 43 Johnson AB, O'Malley BW. Steroid receptor coactivators 1, 2 and 3: critical regulators of nuclear receptor activity and steroid receptor modulator (SRM)-based cancer therapy. *Mol Cell Endocrinol* 2012; **348**: 430–439.
- 44 Oñate SA, Tsai SY, Tsai MJ, O'Malley BW. Sequence and characterization of a coactivator for the steroid hormone receptor superfamily. *Science* 1995; **270**: 1354–1357.
- 45 McInerney EM, Tsai MJ, O'Malley BW, Katzenellenbogen BS. Analysis of estrogen receptor transcriptional enhancement by a nuclear hormone receptor coactivator. *Proc Natl Acad Sci USA* 1996; **93**: 10069–10073.
- 46 Tanenbaum DM, Wang Y, Williams SP, Sigler PB. Crystallographic comparison of the estrogen and progesterone receptor's ligand binding domains. *Proc Natl Acad Sci USA* 1998; **95**: 5998–6003.
- 47 Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL. The structural basis of estrogen receptor/coactivator recognition and the antagonism of this interaction by tamoxifen. *Cell* 1998; **95**: 927–937.
- 48 Oñate SA, Boonyaratanakornkit V, Spencer TE, Tsai SY, Tsai MJ, Edwards DP, O'Malley BW. The steroid receptor coactivator-1 contains multiple receptor interacting and activation domains that cooperatively enhance the activation function 1 (AF1) and AF2 domains of steroid receptors. *J Biol Chem* 1998; **273**: 12101–12108.
- 49 Webb P, Nguyen P, Shinsako J, Anderson C, Feng W, Nguyen MP, Chen D, Huang SM, Subramanian S, McKinerney E, Katzenellenbogen BS, Stallcup MR, Kushner PJ. Estrogen receptor activation function 1 works by binding p160 coactivator proteins. *Mol Endocrinol* 1998; **12**: 1605–1618.
- 50 Dutertre M, Smith CL. Ligand-independent interactions of p160/steroid receptor coactivators and CREB-Binding Protein (CBP) with estrogen receptor- α : regulation by phosphorylation sites in the A/B region depends on other receptor domains. *Mol Endocrinol* 2003; **17**: 1296–1314.
- 51 Torchia J, Rose DW, Inostroza J, Kamei Y, Westin S, Glass CK, Rosenfeld MG. The transcriptional co-activator p/CIP binds CBP and mediates nuclear-receptor function. *Nature* 1997; **387**: 677–684.
- 52 Misiti S, Schomburg L, Yen PM, Chin WW. Expression and hormonal regulation of coactivator and corepressor genes. *Endocrinology* 1998; **139**: 2493–2500.
- 53 Martinez de Arrieta C, Koibuchi N, Chin WW. Coactivator and corepressor gene expression in rat cerebellum during postnatal development and the effect of altered thyroid status. *Endocrinology* 2000; **141**: 1693–1698.
- 54 Meijer OC, Steenbergen PJ, de Kloet ER. Differential expression and regional distribution of steroid receptor coactivators SRC-1 and SRC-2 in brain and pituitary. *Endocrinology* 2000; **141**: 2192–2199.
- 55 Auger AP, Tetel MJ, McCarthy MM. Steroid receptor coactivator-1 mediates the development of sex specific brain morphology and behavior. *Proc Natl Acad Sci USA* 2000; **97**: 7551–7555.
- 56 Molenda HA, Griffin AL, Auger AP, McCarthy MM, Tetel MJ. Nuclear receptor coactivators modulate hormone-dependent gene expression in brain and female reproductive behavior in rats. *Endocrinology* 2002; **143**: 436–444.

- 57 Ogawa H, Nishi M, Kawata M. Localization of nuclear coactivators p300 and steroid receptor coactivator 1 in the rat hippocampus. *Brain Res* 2001; **890**: 197–202.
- 58 Nishihara E, Yoshida-Kimoya H, Chan C, Liao L, Davis RL, O'Malley BW, Xu J. SRC-1 null mice exhibit moderate motor dysfunction and delayed development of cerebellar Purkinje cells. *J Neurosci* 2003; **23**: 213–222.
- 59 Setiawan E, Owen D, McCabe L, Kostaki A, Andrews MH, Matthews SG. Glucocorticoids do not alter developmental expression of hippocampal or pituitary steroid receptor coactivator-1 and -2 in the late gestation fetal guinea pig. *Endocrinology* 2004; **145**: 3796–3803.
- 60 Zalachoras I, Grootaers G, van Weert LT, Aubert Y, de Kreijl SR, Datson NA, van Roon-Mom WM, Aartsma-Rus A, Meijer OC. Antisense-mediated isoform switching of steroid receptor coactivator-1 in the central nucleus of the amygdala of the mouse brain. *BMC Neurosci* 2013; **14**: 5.
- 61 Zhang D, Guo Q, Bian C, Zhang J, Lin S, Su B. Alterations of steroid receptor coactivator-1 (SRC-1) immunoreactivities in specific brain regions of young and middle-aged female Sprague-Dawley rats. *Brain Res* 2011; **1382**: 88–97.
- 62 Yore MA, Im D, Webb LK, Zhao Y, Chadwick JG, Molenda-Figueira HA, Haidacher SJ, Denner L, Tetel MJ. Steroid receptor coactivator-2 expression in brain and physical associations with steroid receptors. *Neuroscience* 2010; **169**: 1017–1028.
- 63 McGinnis MY, Lumia AR, Tetel MJ, Molenda-Figueira HA, Possidente B. Effects of anabolic androgenic steroids on the development and expression of running wheel activity and circadian rhythms in male rats. *Physiol Behav* 2007; **92**: 1010–1018.
- 64 Apostolakis EM, Ramamurthy M, Zhou D, Onate S, O'Malley B. Acute disruption of select steroid receptor coactivators prevents reproductive behavior in rats and unmasks genetic adaptation in knockout mice. *Mol Endocrinol* 2002; **16**: 1511–1523.
- 65 Nottebohm F, Arnold AP. Sexual dimorphism in vocal control areas of the songbird brain. *Science* 1976; **194**: 211–213.
- 66 Jacobs EC, Arnold AP, Campagnoni AT. Developmental regulation of the distribution of aromatase- and estrogen-receptor- mRNA-expressing cells in the zebra finch brain. *Dev Neurosci* 1999; **21**: 453–472.
- 67 Perlman WR, Arnold AP. Expression of estrogen receptor and aromatase mRNAs in embryonic and posthatch zebra finch brain. *J Neurobiol* 2003; **55**: 204–219.
- 68 Saldanha CJ, Coomaringam L. Overlap and co-expression of estrogen synthetic and responsive neurons in the songbird brain – a double-label immunocytochemical study. *Gen Comp Endocrinol* 2005; **141**: 66–75.
- 69 Gurney ME, Konishi M. Hormone-induced sexual differentiation of brain and behavior in zebra finches. *Science* 1980; **208**: 1380–1383.
- 70 Simpson HB, Vicario DS. Early estrogen treatment of female zebra finches masculinizes the brain pathway for learned vocalizations. *J Neurobiol* 1991; **22**: 777–793.
- 71 Grisham W, Mathews GA, Arnold AP. Local intracerebral implants of estrogen masculinize some aspects of the zebra finch song system. *J Neurobiol* 1994; **25**: 185–196.
- 72 Adkins-Regan E, Mansukhani V, Seiwert C, Thompson R. Sexual differentiation of brain and behavior in the zebra finch: critical periods for effects of early estrogen treatment. *J Neurobiol* 1994; **25**: 865–877.
- 73 Fusani L, Metzendorf R, Hutchison JB, Gahr M. Aromatase inhibition affects testosterone-induced masculinization of song and the neural song system in female canaries. *J Neurobiol* 2003; **54**: 370–379.
- 74 Charlier TD, Ball GF, Balthazart J. Rapid action on neuroplasticity precedes behavioral activation by testosterone. *Horm Behav* 2008; **54**: 488–495.
- 75 Serebinski AL, Ball GF, Balthazart J, Charlier TD. Specific activation of estrogen receptor alpha and beta enhances male sexual behavior and neuroplasticity in male Japanese quail. *PLoS ONE* 2011; **6**: e18627.
- 76 Balthazart J, Surlémont C. Copulatory behavior is controlled by the sexually dimorphic nucleus of the quail POA. *Brain Res Bull* 1990; **25**: 7–14.
- 77 Charlier TD, Serebinski AL, Niessen NA, Balthazart J. Modulation of testosterone-dependent male sexual behavior and the associated neuroplasticity. *Gen Comp Endocrinol* 2013Mar 20. pii: S0016-6480(13)00111-1. doi: 10.1016/j.ygcen.2013.03.003. [Epub ahead of print]
- 78 Balthazart J, Surlémont C. Androgen and estrogen action in the preoptic area and activation of copulatory behavior in quail. *Physiol Behav* 1990; **48**: 599–609.
- 79 Charlier TD, Balthazart J, Ball GF. Sex differences in the distribution of the steroid coactivator SRC-1 in the song control nuclei of male and female canaries. *Brain Res* 2003; **959**: 263–274.
- 80 Duncan KA, Jimenez P, Carruth LL. Distribution and sexually dimorphic expression of steroid receptor coactivator-1 (SRC-1) in the zebra finch brain. *Gen Comp Endocrinol* 2011; **170**: 408–414.
- 81 Niessen NA, Balthazart J, Ball GF, Charlier TD. Steroid receptor coactivator 2 modulates steroid-dependent male sexual behavior and neuroplasticity in Japanese quail (*Coturnix japonica*). *J Neurochem* 2011; **119**: 579–593.
- 82 MacLusky NJ, McEwen BS. Oestrogen modulates progesterin receptor concentrations in some rat brain regions but not in others. *Nature* 1978; **274**: 276–278.
- 83 Lauber AH, Romano GJ, Pfaff DW. Sex difference in estradiol regulation of progesterin receptor messenger RNA in rat mediobasal hypothalamus as demonstrated by *In situ* hybridization. *Neuroendocrinology* 1991; **53**: 608–613.
- 84 Blaustein JD, Turcotte JC. Estradiol-induced progesterin receptor immunoreactivity is found only in estrogen receptor-immunoreactive cells in guinea pig brain. *Neuroendocrinology* 1989; **49**: 454–461.
- 85 Warembourg M, Jolivet A, Milgrom E. Immunohistochemical evidence of the presence of estrogen and progesterone receptors in the same neurons of the guinea pig hypothalamus and preoptic area. *Brain Res* 1989; **480**: 1–15.
- 86 Scott REM, Wu-Peng XS, Pfaff DW. Regulation and expression of progesterone receptor mRNA isoforms A and B in the male and female rat hypothalamus and pituitary following oestrogen treatment. *J Neuroendocrinol* 2002; **14**: 175–183.
- 87 Brinton RD, Thompson RF, Foy MR, Baudry M, Wang J, Finch CE, Morgan TE, Pike CJ, Mack WJ, Stanczyk FZ, Nilsen J. Progesterone receptors: form and function in brain. *Front Neuroendocrinol* 2008; **29**: 313–339.
- 88 Tetel MJ, Siegal NK, Murphy SD. Cells in behaviourally relevant brain regions coexpress nuclear receptor coactivators and ovarian steroid receptors. *J Neuroendocrinol* 2007; **19**: 262–271.
- 89 Tognoni CM, Chadwick JG Jr, Ackeifi CA, Tetel MJ. Nuclear receptor coactivators are coexpressed with steroid receptors and regulated by estradiol in mouse brain. *Neuroendocrinology* 2011; **94**: 49–57.
- 90 Zhu L, Yang Y, Xu P, Zou F, Yan X, Liao L, Xu J, O'Malley BW, Xu Y. Steroid receptor coactivator-1 mediates estrogenic actions to prevent body weight gain in female mice. *Endocrinology* 2013; **154**: 150–158.
- 91 Xu J, Qiu Y, Demayo FJ, Tsai SY, Tsai MJ, O'Malley BW. Partial hormone resistance in mice with disruption of the steroid receptor coactivator-1 (SRC-1) gene. *Science* 1998; **279**: 1922–1925.
- 92 Weiss RE, Xu J, Ning G, Pohlenz J, O'Malley BW, Refetoff S. Mice deficient in the steroid receptor co-activator 1 (SRC-1) are resistant to thyroid hormone. *EMBO J* 1999; **18**: 1900–1904.
- 93 Bousios S, Karandrea D, Kittas C, Kitraki E. Effects of gender and stress on the regulation of steroid receptor coactivator-1 expression in the rat brain and pituitary. *J Steroid Biochem Mol Biol* 2001; **78**: 401–407.

- 94 Bian C, Zhang D, Guo Q, Cai W, Zhang J. Localization and sex-difference of steroid receptor coactivator-1 immunoreactivities in the brain of adult female and male mice. *Steroids* 2011; **76**: 269–279.
- 95 Mitev YA, Wolf SS, Almeida OF, Patchev VK. Developmental expression profiles and distinct regional estrogen responsiveness suggest a novel role for the steroid receptor coactivator SRC-1 as a discriminative amplifier of estrogen signaling in the rat brain. *FASEB J* 2003; **17**: 518–519.
- 96 Camacho-Arroyo I, Neri-Gomez T, Gonzalez-Arenas A, Guerra-Araiza C. Changes in the content of steroid receptor coactivator-1 and silencing mediator for retinoid and thyroid hormone receptors in the rat brain during the estrous cycle. *J Steroid Biochem Mol Biol* 2005; **94**: 267–272.
- 97 Zhang D, Guo Q, Bian C, Zhang J, Cai W, Su B. Expression of steroid receptor coactivator-1 was regulated by postnatal development but not ovariectomy in the hippocampus of rats. *Dev Neurosci* 2011; **33**: 57–63.
- 98 Maerkel K, Durrer S, Henseler M, Schlumpf M, Lichtensteiger W. Sexually dimorphic gene regulation in brain as a target for endocrine disruptors: developmental exposure of rats to 4-methylbenzylidene camphor. *Toxicol Appl Pharmacol* 2007; **218**: 152–165.
- 99 Faass O, Schlumpf M, Reolon S, Henseler M, Maerkel K, Durrer S, Lichtensteiger W. Female sexual behavior, estrous cycle and gene expression in sexually dimorphic brain regions after pre- and postnatal exposure to endocrine active UV filters. *Neurotoxicology* 2009; **30**: 249–260.
- 100 Tetel MJ, Ungar TC, Hassan B, Bittman EL. Photoperiodic regulation of androgen receptor and steroid receptor coactivator-1 in Siberian hamster brain. *Mol Brain Res* 2004; **131**: 79–87.
- 101 Iannacone EA, Yan AW, Gauger KJ, Dowling ALS, Zoeller RT. Thyroid hormone exerts site-specific effects on SRC-1 and NCoR expression selectively in the neonatal rat brain. *Mol Cell Endocrinol* 2002; **186**: 49–59.
- 102 Ramos HE, Weiss RE. Regulation of nuclear coactivator and corepressor expression in mouse cerebellum by thyroid hormone. *Thyroid* 2006; **16**: 211–216.
- 103 Charlier TD, Ball GF, Balthazart J. Plasticity in the expression of the steroid receptor coactivator-1 in the Japanese quail brain: effect of sex, testosterone, stress and time of the day. *Neuroscience* 2006; **172**: 333–343.
- 104 Kurihara I, Shibata H, Suzuki T, Ando T, Kobayashi S, Hayashi M, Saito I, Saruta T. Expression and regulation of nuclear receptor coactivators in glucocorticoid action. *Mol Cell Endocrinol* 2002; **189**: 181–189.
- 105 Meijer OC, Kalkhoven E, van der LS, Steenbergen PJ, Houtman SH, Dijkmans TF, Pearce D, de Kloet ER. Steroid receptor coactivator-1 splice variants differentially affect corticosteroid receptor signaling. *Endocrinology* 2005; **146**: 1438–1448.
- 106 Bittman EL, Hegarty CM, Layden MQ, Jonassen JA. Influences of photoperiod on sexual behaviour, neuroendocrine steroid receptors and adeno-hypophysial hormone secretion and gene expression in female golden hamsters. *J Mol Endocrinol* 1990; **5**: 15–25.
- 107 Charlier TD. Importance of steroid receptor coactivators in the modulation of steroid action on brain and behavior. *Psychoneuroendocrinology* 2009; **34**: S20–S29.
- 108 Nishihara E, Moriya T, Shinohara K. Expression of steroid receptor coactivator-1 is elevated during neuronal differentiation of murine neural stem cells. *Brain Res* 2007; **1135**: 22–30.
- 109 Han SJ, Lonard DM, O'Malley BW. Multi-modulation of nuclear receptor coactivators through posttranslational modifications. *Trends Endocrinol Metab* 2009; **20**: 8–15.
- 110 Rowan BG, Weigel NL, O'Malley BW. Phosphorylation of steroid receptor coactivator-1. Identification of the phosphorylation sites and phosphorylation through the mitogen-activated protein kinase pathway. *J Biol Chem* 2000; **275**: 4475–4483.
- 111 Rowan BG, Garrison N, Weigel NL, O'Malley BW. 8-Bromo-cyclic AMP induces phosphorylation of two sites in SRC-1 that facilitate ligand-independent activation of the chicken progesterone receptor and are critical for functional cooperation between SRC-1 and CREB binding protein. *Mol Cell Biol* 2000; **20**: 8720–8730.
- 112 Hoang T, Fenne IS, Cook C, Børud B, Bakke M, Lien EA, Mellgren G. cAMP-dependent protein kinase regulates ubiquitin-proteasome-mediated degradation and subcellular localization of the nuclear receptor coactivator GRIP1. *J Biol Chem* 2004; **279**: 49120–49130.
- 113 Narayanan R, Adigun AA, Edwards DP, Weigel NL. Cyclin-dependent kinase activity is required for progesterone receptor function: novel role for cyclin A/Cdk2 as a progesterone receptor coactivator. *Mol Cell Biol* 2005; **25**: 264–277.
- 114 Gianni M, Parrella E, Raska I Jr, Gaillard E, Nigro EA, Gaudon C, Garattini E, Rochette-Egly C. P38MAPK-dependent phosphorylation and degradation of SRC-3/AIB1 and RARalpha-mediated transcription. *EMBO J* 2006; **25**: 739–751.
- 115 Pleim ET, Brown TJ, MacLusky NJ, Etgen AM, Barfield RJ. Dilute estradiol implants and progestin receptor induction in the ventromedial nucleus of the hypothalamus: correlation with receptive behavior in female rats. *Endocrinology* 1989; **124**: 1807–1812.
- 116 Smith CL, Oñate SA, Tsai MJ, O'Malley BW. CREB binding protein acts synergistically with steroid receptor coactivator-1 to enhance steroid receptor-dependent transcription. *Proc Natl Acad Sci USA* 1996; **93**: 8884–8888.
- 117 Tetel MJ, Giangrande PH, Leonhardt SA, McDonnell DP, Edwards DP. Hormone-dependent interaction between the amino- and carboxyl-terminal domains of progesterone receptor *in vitro* and *in vivo*. *Mol Endocrinol* 1999; **13**: 910–924.
- 118 Sharma D, Handa RJ, Uht RM. The ERbeta ligand 5alpha-androstane, 3beta,17beta-diol (3beta-diol) regulates hypothalamic oxytocin (Oxt) gene expression. *Endocrinology* 2012; **153**: 2353–2361.
- 119 Charlier TD, Harada N, Ball GF, Balthazart J. Targeting steroid receptor coactivator-1 expression with locked nucleic acids antisense reveals different thresholds for the hormonal regulation of male sexual behavior in relation to aromatase activity and protein expression. *Behav Brain Res* 2006; **172**: 333–343.
- 120 Hemmerich P, von Mikecz A, Neumann F, Sözeri O, Wolff-Vorbeck G, Zobelein R, Krawinkel U. Structural and functional properties of ribosomal protein L7 from humans and rodents. *Nucleic Acid Res* 1993; **21**: 223–231.
- 121 Jackson TA, Richer JK, Bain DL, Takimoto GS, Tung L, Horwitz KB. The partial agonist activity of antagonist-occupied steroid receptors is controlled by a novel hinge domain-binding coactivator L7/SPA and the corepressors N-CoR or SMRT. *Mol Endocrinol* 1997; **11**: 693–705.
- 122 Berghöfer-Hochheimer Y, Zurek C, Wölfl S, Hemmerich P, Munder T. L7 protein is a coregulator of vitamin D receptor-retinoid X receptor-mediated transactivation. *J Cell Biochem* 1998; **69**: 1–12.
- 123 Duncan KA, Carruth LL. The sexually dimorphic expression of L7/SPA, an estrogen receptor coactivator, in zebra finch telencephalon. *Dev Neurobiol* 2007; **67**: 1852–1866.
- 124 Duncan KA, Carruth LL. The song remains the same: coactivators and sex differences in the songbird brain. *Front Neuroendocrinol* 2011; **32**: 84–94.
- 125 Duncan KA, Jimenez P, Carruth LL. The selective estrogen receptor-alpha coactivator, RPL7, and sexual differentiation of the songbird brain. *Psychoneuroendocrinology* 2009; **34**(Suppl.): S30–S38.
- 126 Gonzalez-Arenas A, Hansberg-Pastor V, Hernandez-Hernandez OT, Gonzalez-Garcia TK, Henderson-Villalpando J, Lemus-Hernandez D, Cruz-Barríos A, Rivas-Suarez M, Camacho-Arroyo I. Estradiol increases

- cell growth in human astrocytoma cell lines through ERalpha activation and its interaction with SRC-1 and SRC-3 coactivators. *Biochim Biophys Acta* 2012; **1823**: 379–386.
- 127 Hernandez-Hernandez OT, Gonzalez-Garcia TK, Camacho-Arroyo I. Progesterone receptor and SRC-1 participate in the regulation of VEGF, EGFR and Cyclin D1 expression in human astrocytoma cell lines. *J Steroid Biochem Mol Biol* 2012; **132**: 127–134.
- 128 Molenda-Figueira HA, Williams CA, Griffin AL, Rutledge EM, Blaustein JD, Tetel MJ. Nuclear receptor coactivators function in estrogen receptor- and progesterone receptor-dependent aspects of sexual behavior in female rats. *Horm Behav* 2006; **50**: 383–392.
- 129 Mazzucco CA, Walker HA, Pawluski JL, Lieblich SE, Galea LA. ERalpha, but not ERbeta, mediates the expression of sexual behavior in the female rat. *Behav Brain Res* 2008; **191**: 111–117.
- 130 Hardy DF, DeBold JF. The relationship between levels of exogenous hormones and the display of lordosis by the female rat. *Horms Behav* 1971; **22**: 87–97.
- 131 Erskine MS. Solicitation behavior in the estrous female rat: a review. *Horms Behav* 1989; **23**: 473–502.
- 132 Charlier TD, Ball GF, Balthazart J. Inhibition of steroid receptor coactivator-1 blocks estrogen and androgen action on male sex behavior and associated brain plasticity. *J Neurosci* 2005; **25**: 906–913.
- 133 Molenda-Figueira HA, Murphy SD, Shea KL, Siegal NK, Zhao Y, Chadwick JG, Denner LA, Tetel MJ. Steroid receptor coactivator-1 from brain physically interacts differentially with steroid receptor subtypes. *Endocrinology* 2008; **149**: 5272–5279.
- 134 Rissman EF. Roles of oestrogen receptors alpha and beta in behavioural neuroendocrinology: beyond Yin/Yang. *J Neuroendocrinol* 2008; **20**: 873–879.
- 135 Zhao Z, Fan L, Fortress AM, Boulware MI, Frick KM. Hippocampal histone acetylation regulates object recognition and the estradiol-induced enhancement of object recognition. *J Neurosci* 2012; **32**: 2344–2351.
- 136 Fugger HN, Foster TC, Gustafsson J, Rissman EF. Novel effects of estradiol and estrogen receptor alpha and beta on cognitive function. *Brain Res* 2000; **883**: 258–264.
- 137 Isgor C, Cecchi M, Kabbaj M, Akil H, Watson SJ. Estrogen receptor beta in the paraventricular nucleus of hypothalamus regulates the neuroendocrine response to stress and is regulated by corticosterone. *Neuroscience* 2003; **121**: 837–845.
- 138 Kraichely DM, Sun J, Katzenellenbogen JA, Katzenellenbogen BS. Conformational changes and coactivator recruitment by novel ligands for estrogen receptor-alpha and estrogen receptor-beta: correlations with biological character and distinct differences among SRC coactivator family members. *Endocrinology* 2000; **141**: 3534–3545.
- 139 Zhao C, Toresson G, Xu L, Koehler KF, Gustafsson JA, Dahlman-Wright K. Mouse estrogen receptor beta isoforms exhibit differences in ligand selectivity and coactivator recruitment. *Biochemistry* 2005; **44**: 7936–7944.
- 140 Cvorc A, Tatomer D, Tee MK, Zogovic T, Harris HA, Leitman DC. Selective estrogen receptor-beta agonists repress transcription of proinflammatory genes. *J Immunol* 2008; **180**: 630–636.
- 141 Routledge EJ, White R, Parker MG, Sumpter JP. Differential effects of xenoestrogens on coactivator recruitment by estrogen receptor (ER) alpha and ERbeta. *J Biol Chem* 2000; **275**: 35986–35993.
- 142 Singh M, Su C. Progesterone and neuroprotection. *Horm Behav* 2013; **63**: 284–290.
- 143 Sandstrom NJ, Williams CL. Memory retention is modulated by acute estradiol and progesterone replacement. *Behav Neurosci* 2001; **115**: 384–393.
- 144 Tung L, Abdel-Hafiz H, Shen T, Harvell DM, Nitao LK, Richer JK, Sartorius CA, Takimoto GS, Horwitz KB. Progesterone receptors (PR)-B and -A regulate transcription by different mechanisms: AF-3 exerts regulatory control over coactivator binding to PR-B. *Mol Endocrinol* 2006; **20**: 2656–2670.
- 145 Giangrande PH, Kimbrel A, Edwards DP, McDonnell DP. The opposing transcriptional activities of the two isoforms of the human progesterone receptor are due to differential cofactor binding. *Mol Cell Biol* 2000; **20**: 3102–3115.
- 146 Wen DX, Xu YF, Mais DE, Goldman ME, McDonnell DP. The A and B isoforms of the human progesterone receptor operate through distinct signaling pathways within target cells. *Mol Cell Biol* 1994; **14**: 8356–8364.
- 147 Hall JM, McDonnell DP, Korach KS. Allosteric regulation of estrogen receptor structure, function, and coactivator recruitment by different estrogen response elements. *Mol Endocrinol* 2002; **16**: 469–486.